



## Conditions of Use for CHATEAU herbicide

It is a violation of federal law to use CHATEAU herbicide in a manner inconsistent with its labeling. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA-registered CHATEAU herbicide label.

### Approved Crops:

- ❖ Fruiting Vegetables: tomato, pepper (bell & non-bell), eggplant, pepino, tomatillo and okra
- ❖ Melons: citron melon, muskmelon (such as cantaloupe, casaba, honeydew, etc. and watermelon).

### General Information:

- ❖ Chateau can only be applied in row middles between raised plastic mulched beds that are at least 4 inches higher than the treated row middle and the mulched bed must have a minimum of a 24-inch bed width.
- ❖ Spray must remain between raised beds and contact no more than the bottom 1 inch of the raised bed.
- ❖ Do not apply after crops are transplanted.
- ❖ Do not apply more than 4 ounces of Chateau per acre during a single application.
- ❖ All applications must be made with shielded or hooded equipment.
- ❖ Injury can occur if soil particles treated with Chateau are blown or splashed onto crop.
- ❖ A rainfall after application but prior to transplanting is required.
- ❖

### Rate:

- ❖ Up to 4 ounces/acre per application.
- ❖

### Timing:

- ❖ Chateau may be applied as a shielded or hooded application to row middles after plastic is laid up to transplanting. Spray must be directed to the row middle and contact no more than the bottom 1 inch of the side of the raised bed. If top of much beds is contacted, severe injury can occur due to foliage contact with treated plastic.

### Weed Control and Tank Mixing

- ❖ Chateau provides preemergence residual control of Broadleaf Weed Species and Grass Weed Species as well as to assist in the postemergence control of emerged weeds. **\*\*Refer to Federal Label for list of weeds controlled.** A registered preemergence grass herbicide may be added for control of additional grassy weeds. For control of emerged weeds, tank-mix Chateau with paraquat, Aim, glyphosate, or other registered burndown herbicide. Refer to tank mix partner label for recommended rates and application parameters.

## **Rotational Restrictions:**

Multi-Crop Plastic Culture: Chateau may only be used in row middles of registered crops at time of application. However, if an unregistered crop is planted back to the same undisturbed plastic mulch after primary crop is removed; there is a rotational restriction of 30 days following original application.

If plastic mulch is removed and soil is re-bedded, follow the following rotational restrictions:

- 1) **Crops:** Cotton, Field Corn, Peanut, Rice, Sorghum, Soybean, Sunflower, Tobacco and Wheat. **Rotation Intervals:** 4 months. **Rate:** Up to 4 oz/A
- 2) **Crops:** Alfalfa, Canola, Sugar Beet and all other crops not listed. **Rotation Intervals:** 6 months if soil is tilled prior to planting. 12 months if no tillage is performed. **Rate:** Up to 4 oz/A.

## **Tank Cleanout**

Refer to Federal label for tank cleanout procedures. A tank cleaner specifically designed for Chateau may be purchased through your local dealer.

The decision to use or not use this product must be made by the user and/or grower on the basis of possible crop injury from CHATEAU herbicide, the severity of weed infestation, the cost of alternate weed controls, and other factors. Because of the risk of crop damage, all such use is at user and/or grower's risk.

---